

THE GALEX UV LUMINOSITY FUNCTION OF THE CLUSTER OF GALAXIES ABELL 1367.

L. CORTESE¹, A. BOSELLI¹, G. GAVAZZI², J. IGLESIAS-PARAMO¹, B. F. MADORE^{3,4}, T. BARLOW⁵, L. BIANCHI⁶, Y.-I. BYUN⁷, J. DONAS¹, K. FORSTER⁵, P. G. FRIEDMAN⁵, T. M. HECKMAN⁸, P. JELINSKY⁹, Y.-W. LEE⁷, R. MALINA¹, D. C. MARTIN⁵, B. MILLIARD¹, P. MORRISSEY⁵, S. NEFF¹⁰, R. M. RICH¹¹, D. SCHIMINOVICH⁵, O. SIEGMUND⁹, T. SMALL⁵, A. S. SZALAY⁸, M. A. TREYER⁵, B. WELSH⁹, T. K. WYDER⁵

Draft version February 2, 2008

ABSTRACT

We present the GALEX NUV (2310Å) and FUV (1530Å) galaxy luminosity functions of the nearby cluster of galaxies A1367 in the magnitude range $-20.3 \leq M_{AB} \leq -13.3$. The luminosity functions are consistent with previous (~ 2 mag shallower) estimates based on the FOCA and FAUST experiments, but display a steeper faint-end slope than the GALEX luminosity function for local field galaxies. Using spectro-photometric optical data we select out star-forming systems from quiescent galaxies and study their separate contributions to the cluster luminosity function. We find that the UV luminosity function of cluster star-forming galaxies is consistent with the field. The difference between the cluster and field LF is entirely due to the contribution at low luminosities ($M_{AB} > -16$ mag) of non star-forming, early-type galaxies that are significantly over dense in clusters.

Subject headings: ultraviolet: galaxies – galaxies:luminosity function, evolution – galaxies: clusters: individual (Abell1367)

1. INTRODUCTION

Recent determinations of the galaxy luminosity function (hereafter LF) at various frequencies, in various environments (i.e. De Propris et al. 2003; Madgwick et al. 2002) and in a number of redshift intervals (i.e. Ilbert et al. 2004) have improved our knowledge of galaxy evolution and the role played by the environment in regulating the star formation activity of galaxies. The optical cluster LF is significantly steeper than that in the field (Trentham et al. 2005). This steepening is due to quiescent galaxies, more frequent at low luminosities in clusters, while the LF of cluster star forming objects is similar to that in the field (De Propris et al. 2003). The causes of this difference might reside in the density-morphology relation (Dressler 1980; Whitmore et al. 1993) and in particular in the overabundance of dwarf ellipticals in rich clusters (Ferguson & Sandage 1991), whose origin is currently debated in the framework of the environmental effects

on galaxy evolution.

The ultraviolet emission UV(~ 2000 Å), being dominated by young stars of intermediate masses ($2 < M < 5 M_{\odot}$, Boselli et al. 2001) represents an appropriate tool to identify and quantify star formation activity. Before the launch of the *Galaxy Evolution Explorer* (GALEX) a few balloon-borne experiments had observed the sky at ultraviolet wavelengths (Smith & Cornett 1982; Lampton et al. 1990; Kodaira et al. 1990). Among them, the FOCA experiment (Milliard et al. 1991) allowed the first determinations of the UV LF of local field galaxies (Treyer et al. 1998; Sullivan et al. 2000) and of nearby clusters (Donas et al. 1991; Andreon 1999; Cortese et al. 2003b). Combining the FOCA and FAUST data Cortese et al. (2003a) determined the first composite LF of nearby clusters. They found no significant differences with the LF in the field. However this early determination was affected by large statistical errors due to the lack of sufficient redshift coverage for UV selected galaxies and to the uncertainty in the UV background counts (Cortese et al. 2003b). GALEX has opened a new era of extragalactic UV astronomy. In particular it provides for the first time precise UV photometry of galaxies over large stretches of the sky (Xu et al. 2005), thus making the background subtraction method more reliable than in the past.

In this letter we present the first determination of the UV (NUV and FUV) LF in a nearby cluster of galaxies based upon GALEX measurements of galaxies brighter than $M_{AB} \sim -13.3$ that extends two magnitudes deeper than previous estimates (Cortese et al. 2003a). Throughout this paper we assume for Abell 1367 a distance modulus $\mu = 34.84$ mag (Sakai et al. 2000), corresponding to $H_0 = 70$ km s⁻¹ Mpc⁻¹.

2. UV DATA

GALEX provides far-ultraviolet (FUV; $\lambda_{\text{eff}} = 1530\text{Å}$, $\Delta\lambda = 400\text{Å}$) and near-ultraviolet (NUV; $\lambda_{\text{eff}} =$

¹ Laboratoire d'Astrophysique de Marseille, BP8, Traverse du Siphon, F-13376 Marseille, France

² Università degli Studi di Milano - Bicocca, P.zza della Scienza 3, 20126 Milano, Italy

³ Observatories of the Carnegie Institution of Washington, 813 Santa Barbara St., Pasadena, CA 91101

⁴ NASA/IPAC Extragalactic Database, California Institute of Technology, Mail Code 100-22, 770 S. Wilson Ave., Pasadena, CA 91125

⁵ California Institute of Technology, MC 405-47, 1200 East California Boulevard, Pasadena, CA 91125

⁶ Center for Astrophysical Sciences, The Johns Hopkins University, 3400 N. Charles St., Baltimore, MD 21218

⁷ Center for Space Astrophysics, Yonsei University, Seoul 120-749, Korea

⁸ Department of Physics and Astronomy, The Johns Hopkins University, Homewood Campus, Baltimore, MD 21218

⁹ Space Sciences Laboratory, University of California at Berkeley, 601 Campbell Hall, Berkeley, CA 94720

¹⁰ Laboratory for Astronomy and Solar Physics, NASA Goddard Space Flight Center, Greenbelt, MD 20771

¹¹ Department of Physics and Astronomy, University of California, Los Angeles, CA 90095




FIG. 1.— The redshift completeness per bin of UV magnitude in Abell 1367.

2310Å, $\Delta\lambda = 1000\text{\AA}$) images with a circular field of view of ~ 0.6 degrees radius. The spatial resolution is ~ 5 arcsec. Details of the GALEX instrument can be found in Martin et al. (2005) and Morrissey et al. (2005). The data analyzed in this paper consist of two GALEX pointings of the Abell cluster 1367, with a mean exposure time of 1460s, centered at R.A.(J2000)=11:43:41.34 Dec(J2000)=+20:11:24.0 (e.g. offset to the north of the cluster to avoid a star bright enough to threaten the detector). Sources were detected and measured using SExtractor (Bertin & Arnouts 1996). The 100% completeness limit is $m_{AB} \sim 21.5$ both in FUV and NUV (Xu et al. 2005). As the NUV images are significantly deeper than the FUV, sources were selected and their parameters determined in the NUV. FUV parameters were extracted in the same apertures. We used a larger SExtractor deblending parameter compared to the standard GALEX pipeline, providing reliable MAG_{AUTO} also for very extended sources. The calibration uncertainty of the NUV and FUV magnitudes is $\sim 10\%$ (Morrissey et al. 2005). Magnitudes are corrected for Galactic extinction using the Schlegel et al. (1998) reddening map and the Galactic extinction curve of Cardelli et al. (1989). The applied extinction corrections are of 0.18 and 0.17 mag for the NUV and FUV bands respectively. To avoid artifacts present at the edge of the field, we considered only the central 0.58 deg radius from the field center. A reliable star/galaxy discrimination was achieved by matching the GALEX catalog against the deepest optical catalogs available for A1367 ($B < 22.5$ mag and $r' < 21$ mag; Iglesias-Páramo et al. 2003), using a search radius of 6 arcsec, as adopted by Wyder et al. (2005) for the estimate of the GALEX local field LF. The optical catalogs do not include a negligible part (~ 0.09 square degrees) of the GALEX field. A total number of 292 galaxies in the FUV and of 480 galaxies in NUV with $m_{AB} \leq 21.5$ are detected in the ~ 0.96 square degrees field ($\sim 2.5\text{Mpc}^2$) analyzed in this work.

3. THE LUMINOSITY FUNCTION

TABLE 1
BEST FITTING PARAMETERS.

Band	Sample	Schechter Parameters	
		M^*	α
NUV	A1367	-19.77 ± 0.42	-1.64 ± 0.21
NUV	Field	-18.23 ± 0.11	-1.16 ± 0.07
FUV	A1367	-19.86 ± 0.50	-1.56 ± 0.19
FUV	Field	-18.04 ± 0.11	-1.22 ± 0.07
UV(2000Å)	Composite cluster	-18.79 ± 0.40	-1.50 ± 0.10

The determination of the cluster LF requires a reliable estimate of the contribution from background/foreground objects to the UV counts. This can be accurately achieved for $m_{AB} \leq 18.5$, since at this limit our redshift completeness is $\sim 90\%$ (Cortese et al. 2003b, 2004; see Fig. 1).

The redshift completeness drops rapidly at magnitudes fainter than $m_{AB} \sim 18.5$, thus requiring the contamination to be estimated statistically. Two methods are usually applied for the computation of cluster LFs. The first one is based on the statistical subtraction of field galaxies, per bin of UV magnitude, that are expected to be randomly projected onto the cluster area, as derived by Xu et al. (2005). Alternatively, the *completeness corrected* method proposed by De Propris et al. (2003) is to be preferred when the field counts have large uncertainties. It is based on the assumption that the UV spectroscopic sample (e.g. membership confirmed spectroscopically) is ‘representative’ of the entire cluster, i.e. the fraction of galaxies that are cluster members is the same in the (incomplete) spectroscopic sample as in the (complete) photometric one. For each magnitude bin i we count the number of cluster members N_M (i.e. galaxies with velocity in the range $4000 < V < 10000 \text{ km s}^{-1}$; Cortese et al. 2004), the number of galaxies with a measured recessional velocity N_Z and the total number of galaxies N_T . The ratio N_Z/N_T , corresponding to the redshift completeness in each magnitude bin is shown in Fig.1. The completeness-corrected number of cluster members in each bin is $N_i = (N_M \times N_T)/N_Z$.

N_T is a Poisson variable, and N_M is a binomial variable (the number of successes in N_Z trials with probability N_M/N_Z). Therefore the errors associated with N_i are given by $(\delta^2 N_i / N_i^2) = (1/N_T) + (1/N_M) - (1/N_Z)$. The NUV and FUV LFs using both methods (see Fig 2) are in good agreement for $M_{AB} \geq -14.3$. In the last bin the two methods are inconsistent as the *completeness corrected* method predicts a higher slope than the statistical background subtraction. This disagreement is likely due to the severe redshift incompleteness for $M_{AB} \geq -14.3$. In any case we take the weighted mean of the two determinations.

Due to the small number of galaxies populating the high luminosity bins (i.e. only three objects brighter than $M_{AB} \sim -18.3$), the LFs are not well fitted with a Schechter function (Schechter 1976): the best-fit M_* turns out to be brighter than the brightest observed galaxy. For this reason we first determine the faint-end ($-18.3 \leq M_{AB} \leq -13.3$) slope in each band, fitting the LFs with a power law ($\Phi(M) = c 10^{kM}$) by minimizing




FIG. 2.— The GALEX NUV (left) and FUV (right) LF for Abell 1367. Open dots are obtained using the subtraction of field counts obtained by Xu et al. (2005); filled dots are obtained using the *completeness corrected* method. The solid line represents the best Schechter fit. The dotted line shows the composite nearby clusters 2000 Å LF by (Cortese et al. 2003a). The dashed line represents the GALEX local field LF (Wyder et al. 2005), normalized in order to match the cluster LF at $M_{AB} \sim -17.80$.




FIG. 3.— The NUV (left) and FUV (right) bi-variate LF of A1367. Star-forming and quiescent galaxies are indicated with empty triangles and filled squares respectively. The dashed line represents the GALEX local field LF (Wyder et al. 2005), normalized as in Fig.2

χ^2 . The α parameter of the Schechter function can be derived from k using the relation $\alpha = -(k/0.4 + 1)$. Then we fit the LFs with a Schechter function, keeping α fixed to the value previously obtained. This is not the canonical Schechter fit, but it provides a more realistic set of parameters than using a three-free-parameter fit. The best fit parameters and their errors are listed in Table.1. In order to separate the contribution to the LF of star-forming from quiescent galaxies, we divide the sample into two classes. Using $H\alpha$ imaging data (Iglesias-Páramo et al. 2002; Gavazzi et al. 1998, 2002, 2003) and optical spectroscopy (Cortese et al. 2003b, 2004) we can discriminate between star-forming ($EW(H\alpha) > 0 \text{ \AA}$) and quiescent ($EW(H\alpha) = 0 \text{ \AA}$) objects. Unfortunately neither UV field counts for different morphological types nor a measure of $EW(H\alpha)$ for all the UV selected galaxies are available. Thus we can only

apply the *completeness corrected* method to determine the bi-variate LFs. We assume that in each bin of magnitude the fraction of star-forming and quiescent cluster members is the same in the (incomplete) spectroscopic sample as in the (complete) photometric sample. The bi-variate LFs derived by this method are shown in Fig.3.

4. DISCUSSION

As shown in Fig.2, the GALEX LFs have a shape consistent to the composite LF of nearby clusters as constructed by Cortese et al. (2003a). Conversely, whatever fitting procedure one adopts, they show a steeper faint-end slope and a brighter M^* than the GALEX field LF recently determined by Wyder et al. (2005). The brighter M^* observed in Abell1367 is probably to be ascribed to the particular galaxy population of this cluster. In fact Abell 1367 is a young cluster of galaxies com-




FIG. 4.— The FUV-NUV color magnitude relation for confirmed members of A1367. Symbols are as in Fig.3

posed of at least four dynamical units at the early stage of a multiple merging event (Cortese et al. 2004). Some galaxies have their star formation enhanced due to the interaction with the cluster environment, and it is this population that is responsible for the bright M^* observed in this cluster.

Conversely the high faint-end slope observed in this cluster is due to the significant contribution of non star-forming systems at faint UV magnitudes. In fact, as shown in Fig.3, star-forming galaxies dominate the UV

¹ The bi-variate LF's cannot be compared with the ones computed by Treyer et al. (2005) for the field, since their samples do

not contain ellipticals but only spiral galaxies.

LF for $M_{AB} \leq -17$ mag, as Donas et al. (1991) concluded for the first time. For $M_{AB} \geq -16$ mag however, the number of red galaxies increases very rapidly¹. This result is consistent with an UV LF constructed starting from the r' LF computed by Iglesias-Páramo et al. (2003): assuming a mean color $NUV - r' \sim 1$ mag and $NUV - r' \sim 5$ for star-forming and quiescent galaxies respectively, we are able to reproduce the contribution, at low UV luminosities, of elliptical galaxies. We conclude that a significant fraction of the low luminosity UV emission comes from extreme Horizontal Branch stars and their progenitors (O'Connell 1999). This conclusion is reinforced by the FUV-NUV color magnitude relation (computed only for confirmed cluster members) shown in Fig.4. The star-forming objects dominate at high UV luminosities while the quiescent systems contribute more at faint magnitudes. Their mean FUV-NUV color is ~ 1.5 mag thus they influence the LF at higher luminosities in the NUV than in the FUV (see Fig.3). Finally, the LF's of cluster star-forming systems have a faint-end slope ($\alpha \sim -1.25 \pm 0.2$) consistent within the statistical uncertainties with the GALEX field LF. This result is expected since in the field the fraction of quiescent systems is significantly lower than that of star forming objects (Dressler 1980; Whitmore et al. 1993), thus their contribution to the LF is negligible.

GALEX (Galaxy Evolution Explorer) is a NASA Small Explorer, launched in April 2003. We gratefully acknowledge NASA's support for construction, operation, and science analysis for the GALEX mission, developed in co-operation with the Centre National d'Etudes Spatiales of France and the Korean Ministry of Science and Technology. This research has made extensive use of the GOLD-Mine Database.

REFERENCES

- Andreon, S. 1999, *A&A*, 351, 65
 Bertin, E. & Arnouts, S. 1996, *A&AS*, 117, 393
 Boselli, A., Gavazzi, G., Donas, J., & Scodeggio, M. 2001, *AJ*, 121, 753
 Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245
 Cortese, L., Gavazzi, G., Boselli, A., Iglesias-Páramo, J., & Carrasco, L. 2004, *A&A*, 425, 429
 Cortese, L., Gavazzi, G., Boselli, A., et al. 2003a, *A&A*, 410, L25
 Cortese, L., Gavazzi, G., Iglesias-Páramo, J., Boselli, A., & Carrasco, L. 2003b, *A&A*, 401, 471
 De Propris, R., Colless, M., Driver, S. P., et al. 2003, *MNRAS*, 342, 725
 Donas, J., Milliard, B., & Laget, M. 1991, *A&A*, 252, 487
 Dressler, A. 1980, *ApJ*, 236, 351
 Ferguson, H. C. & Sandage, A. 1991, *AJ*, 101, 765
 Gavazzi, G., Boselli, A., Donati, A., Franzetti, P., & Scodeggio, M. 2003, *A&A*, 400, 451
 Gavazzi, G., Boselli, A., Pedotti, P., Gallazzi, A., & Carrasco, L. 2002, *A&A*, 386, 114
 Gavazzi, G., Catinella, B., Carrasco, L., Boselli, A., & Contursi, A. 1998, *AJ*, 115, 1745
 Iglesias-Páramo, J., Boselli, A., Cortese, L., Vilchez, J. M., & Gavazzi, G. 2002, *A&A*, 384, 383
 Iglesias-Páramo, J., Boselli, A., Gavazzi, G., Cortese, L., & Vilchez, J. M. 2003, *A&A*, 397, 421
 Ilbert, O., Tresse, L., Zucca, E., et al. 2004, *astro-ph/0409134*
 Kodaira, K., Watanabe, T., Onaka, T., & Tanaka, W. 1990, *ApJ*, 363, 422
 Lampton, M., Deharveng, J. M., & Bowyer, S. 1990, in *IAU Symp.* 139, 449
 Madgwick, D. S., Lahav, O., Baldry, I. K., et al. 2002, *MNRAS*, 333, 133
 Martin, D. C., Fanson, J., Schiminovich, D., et al. 2005, *ApJL*, *Galax special issue (astro-ph/0411302)*
 Milliard, B., Donas, J., & Laget, M. 1991, *Advances in Space Research*, 11, 135
 Morrissey, P., Schiminovich, D., Barlow, T. A., et al. 2005, *ApJL*, *Galax special issue (astro-ph/0411310)*
 O'Connell, R. W. 1999, *ARA&A*, 37, 603
 Sakai, S., Mould, J. R., Hughes, S. M. G., et al. 2000, *ApJ*, 529, 698
 Schechter, P. 1976, *ApJ*, 203, 297
 Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
 Smith, A. M. & Cornett, R. H. 1982, *ApJ*, 261, 1
 Sullivan, M., Treyer, M. A., Ellis, R. S., et al. 2000, *MNRAS*, 312, 442
 Trentham, N., Sampson, L. & Banerji, M. 2005, *MNRAS*, in press (*astro-ph/0412124*)
 Treyer, M., Wyder, T. K., Schiminovich, D., et al. 2005, *ApJL*, *Galax special issue (astro-ph/0411308)*
 Treyer, M. A., Ellis, R. S., Milliard, B., Donas, J., & Bridges, T. J. 1998, *MNRAS*, 300, 303
 Whitmore, B. C., Gilmore, D. M., & Jones, C. 1993, *ApJ*, 407, 489
 Wyder, T. K., Treyer, M. A., Milliard, B., et al. 2005, *ApJL*, *Galax special issue (astro-ph/0411364)*
 Xu, C. G., Donas, J., Arnouts, S., et al. 2005, *ApJL*, *Galax special issue (astro-ph/0411317)*